

26.11.2024

ADM

①

L3

WIL3

The symmetric group S_n Let $n \in \mathbb{Z}_{>0}$.The vector space of $n \times n$ matrices $M_n(\mathbb{C})$ has \mathbb{C} -basis $\{E_{ij} \mid i, j \in \{1, \dots, n\}\}$ where E_{ij} is the matrix with 1 in the (i, j) -entry and 0 elsewhere.A permutation of n is $w \in M_{n \times n}(\mathbb{C})$ such that

(a) There is exactly one nonzero entry in each row and each column.

(b) The nonzero entries are 1.

The symmetric group is

$$S_n = \left\{ w \in M_{n \times n}(\mathbb{C}) \mid w \text{ is a permutation of } \{1, \dots, n\} \right\}$$

with matrix multiplication.

Identify a permutation $w \in M_{n \times n}(\mathbb{C})$ with a bijection $w: \{1, \dots, n\} \rightarrow \{1, \dots, n\}$ by

$$w(i) = j \quad \text{if} \quad w_{ij} = 1$$

where w_{ij} is the (i, j) -entry of the matrix w .

26.11.2024 (2)

the transpositions, or reflections, ADM L3 W/L3
 in S_n are

$$s_{ij} = 1 + E_{ij} + E_{ji} - E_{ii} - E_{jj}, \quad \text{for } i, j \in \{1, \dots, n\} \\ \text{with } i \neq j.$$

the simple transpositions are

$$s_1 = s_{12}, \quad s_2 = s_{23}, \quad \dots, \quad s_{n-1} = s_{n-1, n}.$$

The general linear group is

$$GL_n(\mathbb{C}) = \left\{ A \in M_n(\mathbb{C}) \mid \text{there exists } A^{-1} \in M_n(\mathbb{C}) \right. \\ \left. \text{with } AA^{-1} = I \text{ and } A^{-1}A = I \right\}$$

with matrix multiplication

Proposition The maps

$$GL_n(\mathbb{C}) \times GL_m(\mathbb{C}) \longrightarrow GL_{n+m}(\mathbb{C}) \\ (A, B) \longmapsto \left(\begin{array}{c|c} A & 0 \\ \hline 0 & B \end{array} \right) = A \oplus B$$

and

$$S_n \times S_m \longrightarrow S_{n+m} \\ (v, w) \longmapsto \left(\begin{array}{c|c} v & 0 \\ \hline 0 & w \end{array} \right) = v \times w$$

are injective group homomorphisms.

26.11.2024 (3)
ADM. L3
W/L3

Let $\gamma_1 = E_{11} \in S_1$ and

$$\gamma_k = E_{12} + E_{23} + \dots + E_{k-1,k} + E_{k1} \in S_k$$

for $k \in \mathbb{Z}_{>1}$. For $\mu_1, \dots, \mu_\ell \in \mathbb{Z}_{>0}$ let

$$\gamma_\mu = \gamma_{\mu_1} \times \dots \times \gamma_{\mu_\ell} \in S_{\mu_1} \times \dots \times S_{\mu_\ell} \subseteq S_{\mu_1 + \dots + \mu_\ell}.$$

A Coxeter element of S_n is an element of the conjugacy class of γ_n in S_n .

Let $[\gamma_\mu]$ denote the conjugacy class of γ_μ in $S_{\mu_1 + \dots + \mu_\ell}$.

A partition of n is $\lambda = (\lambda_1, \dots, \lambda_\ell)$ such that

$$\lambda_1, \dots, \lambda_\ell \in \mathbb{Z}_{>0} \text{ and } \lambda_1 \geq \dots \geq \lambda_\ell$$

$$\text{and } \lambda_1 + \dots + \lambda_\ell = n.$$

Theorem (a) The map

$$\left\{ \begin{array}{l} \text{partitions} \\ \text{of } n \end{array} \right\} \longleftrightarrow \left\{ \begin{array}{l} \text{conjugacy classes} \\ \text{of } S_n \end{array} \right\}$$

$$\lambda \longmapsto [\gamma_\lambda]$$

is a bijection

26.11.2024 (4)


ADP L3 W/L3

(b) $\text{Card}(\Sigma \gamma_\lambda) = \frac{n!}{z_\lambda}$, where

$z_\lambda = (1^{m_1} 2^{m_2} \dots) (m_1! m_2! \dots)$ if $\lambda = (1^{m_1} 2^{m_2} \dots)$

so that m_i is the number of parts of size i in λ .

Proof



and if

$\gamma_\lambda = \gamma_1 \times \gamma_1 \times \gamma_1 \times \gamma_1 \times \gamma_2 \times \gamma_2 \times \gamma_2 \times \gamma_2 \times \gamma_3 \times \gamma_4 \times \gamma_4$

then

$\text{Card}(\text{stab}(\gamma_\lambda)) = 4! \cdot 1! \cdot 1! \cdot 1! \cdot 1! \cdot 3! \cdot 2 \cdot 2 \cdot 2 \cdot 3 \cdot 2! \cdot 4 \cdot 4$
 $= 4! \cdot 1^4 \cdot 3! \cdot 2^3 \cdot 1! \cdot 3! \cdot 2! \cdot 4^2$

so that

$\text{Card}(\Sigma \gamma_\lambda) = \frac{\text{Card}(S_n)}{\text{Card}(\text{stab}(\gamma_\lambda))} = \frac{n!}{z_\lambda}$