

1.6 The binomial theorem

Let $k \in \mathbb{Z}_{\geq 0}$. Define k factorial by

$$0! = 1 \quad \text{and} \quad k! = k \cdot (k-1) \cdots 3 \cdot 2 \cdot 1 \text{ if } k \in \mathbb{Z}_{>0}.$$

Let $n, k \in \mathbb{Z}_{\geq 0}$ with $k \leq n$. Define

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}.$$

Theorem 1.1. Let $n, k \in \mathbb{Z}_{\geq 0}$ with $k \leq n$.

(a) Let S be a set with cardinality n . Then

$\binom{n}{k}$ is the number of subsets of S with cardinality k .

(b) $\binom{n}{k}$ is the coefficient of $x^{n-k}y^k$ in $(x+y)^n$.

(c) If $k \in \{1, \dots, n-1\}$ then

$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}, \quad \text{and} \quad \binom{n}{0} = 1 \quad \text{and} \quad \binom{n}{n} = 1.$$

This theorem says that the table of numbers

$$\begin{array}{ccccccccccccc} & & & & \binom{0}{0} & & & & & & & & \\ & & & & \binom{1}{0} & \binom{1}{1} & & & & & & & \\ & & & & \binom{2}{0} & \binom{2}{1} & \binom{2}{2} & & & & & & \\ & & & & \binom{3}{0} & \binom{3}{1} & \binom{3}{2} & \binom{3}{3} & & & & & \\ & & & & \binom{4}{0} & \binom{4}{1} & \binom{4}{2} & \binom{4}{3} & \binom{4}{4} & & & & \\ \binom{5}{0} & \binom{5}{1} & \binom{5}{2} & \binom{5}{3} & \binom{5}{4} & \binom{5}{5} & & & & & & & \\ \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & & & & & & & \end{array}$$

are the numbers in **Pascal's triangle**

$$\begin{array}{ccccccccccccc} & & & & 1 & & & & & & & & & \\ & & & & 1 & 1 & 1 & & & & & & & \\ & & & & 1 & 2 & 1 & 1 & & & & & & \\ & & & & 1 & 3 & 3 & 1 & 1 & & & & & \\ & & & & 1 & 4 & 6 & 4 & 1 & 1 & & & & \\ 1 & 5 & 10 & 10 & 5 & 1 & & & & & & & & \\ \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & & & & & & & & \end{array}$$

and that

$$\begin{aligned} (x+y)^0 &= 1, \\ (x+y)^1 &= x+y, \\ (x+y)^2 &= x^2 + 2xy + y^2, \\ (x+y)^3 &= x^3 + 3x^2y + 3xy^2 + y^3, \\ (x+y)^4 &= x^4 + 4x^3y + 6x^2y^2 + 4xy^3 + y^4, \\ (x+y)^5 &= x^5 + 5x^4y + 10x^3y^2 + 10x^2y^3 + 5xy^4 + y^5, \\ \vdots & \vdots \end{aligned}$$